

FLOER THEORY AND TOPOLOGY OF $Diff(S^2)$

YASHA SAVELYEV

ABSTRACT. We say that a fixed point of a diffeomorphism is non-degenerate if 1 not an eigenvalue of the linearization at the fixed point. We use Floer theory to study some aspects of the topology of non-degenerate area preserving diffeomorphism group of the 2-sphere.

1. INTRODUCTION

The group of orientation preserving diffeomorphisms of S^2 has been shown to deformation retract to $SO(3)$ in the classical work of Smale [8]. However it seems we are still very far from completely understanding some finer aspects of its topology. Consider for example the following elementary question.

Question 1.1. *Let $Diff^2(S^2)$ denote the space of orientation preserving diffeomorphisms of S^2 , with exactly 2, prescribed non-degenerate fixed points (and no others). Where non-degenerate means that 1 is not an eigenvalue of the linearization at the fixed point. Is this space contractable?*

We expect that the answer is yes, since $Diff^2(S^2) \cap SO(3)$ is just the space $SO^2(3)$ of rotations about a fixed axis, not containing the identity map, and so forms a contractible space. And so the intuition behind Smale's, and or somewhat related Moser's argument is indicative that there should be a deformation retraction of $Diff^2(S^2) \cap SO(3)$ onto $SO^2(3)$. The tricky point in directly applying Smale's argument (for instance) is that in constructing the deformation retraction one must be careful not to introduce additional fixed points.

While we don't address the question above in this note, we shall look at some very closely related questions (in particular involving versions of the space $Diff^2(S^2)$ above) using Hamiltonian Floer theory.

Let

$$\text{Ham}^{ndg}(S^2) \subset \text{Ham}(S^2, \omega) = \text{DiffVol}(S^2)$$

denote the subspace of area preserving diffeomorphisms ϕ of S^2 with isolated, non-degenerate fixed points. We may further split $\text{Ham}^{ndg}(S^2)$ into components $\text{Ham}(S^2, 2k)$ with $2k \geq 2$, denoting the number of fixed points. Note that these components are not connected, and their topological structure seems to be completely unknown. We note however that there are natural fibrations

$$\text{Ham}(S^2, 2k) \rightarrow C_{2k}(S^2),$$

with the latter denoting the configuration space of unordered points on S^2 , where points are not allowed to collide. While much seems to be known about the topology of the spaces $C_{2k}(S^2)$, see for example McDuff [5], the fiber may be tricky to

Key words and phrases. Floer theory, configuration spaces of points, positivity of intersections, Cerf theory.

understand for higher k , although for $k = 1$ the fiber is the analogue of the space $\text{Diff}^2(S^2)$, and we expect that it is contractible.

Theorem 1.2. *For $i : \text{Ham}^{\text{ndg}}(S^2) \rightarrow \text{Ham}(S^2, \omega)$ the inclusion map, and for $j : B \hookrightarrow \text{Ham}^{\text{ndg}}(S^2)$ a continuous map with B having the homotopy type of a CW complex, s.t. the induced map $i \circ j : B \hookrightarrow \text{Ham}(S^2, \omega)$ is vanishing on π_1 , $i \circ j$ is null-homotopic. In particular i vanishes on higher homotopy groups.*

Our proof involves elliptic techniques in symplectic geometry, and Floer theory in particular, but in a rather transparent way. Note that the above corollary fails if we partially compactify the space $\text{Ham}^{\text{ndg}}(S^2)$ by adding id . Since the natural Lie group homomorphism $S^3 \rightarrow \text{Ham}(S^2, \omega)$ would lie in the corresponding space, and represents the generator.

Remark 1.3. *A technically identical argument but dealing with Hamiltonian fibrations over strips/discs, rather than the cylinder/ S^2 , gives an analogue of the above theorem for the space $\text{Lag}(S^2)$ of Lagrangian submanifolds in S^2 . Hamiltonian isotopic to equator, i.e. simple closed smooth unparametrized loops partitioning S^2 into a pair of regions with equal area. The basic ingredient for this is that a Hamiltonian loop of Lagrangian submanifolds, that is a loop obtained by a Hamiltonian flow, gives rise to a sub-fibration of $M \times D^2$ over the boundary ∂D^2 with fiber $L_\theta \subset M_\theta$ over θ the element of the loop. We may then analogously study moduli spaces of holomorphic disks with boundary on this Lagrangian sub-fibration, for suitable almost complex structures, see for instance [1] for a similar story, or [3] for the monotone case.*

Claim 1.4. *For $i : \text{Lag}^{\text{ndg}}(S^2) \rightarrow \text{Lag}(S^2)$ the inclusion map, and for $j : B \rightarrow \text{Lag}^{\text{ndg}}(S^2)$ a map of a CW complex, s.t. the induced inclusion map $i \circ j : B \hookrightarrow \text{Lag}(S^2)$ is vanishing on π_1 , $i \circ j$ is null-homotopic. Where the superscript ndg means we take the subspace of those Lagrangians intersecting the standard equator at isolated points.*

2. SOME PRELIMINARIES AND NOTATION

2.1. The group of Hamiltonian symplectomorphisms. Given a smooth function $H : M^{2n} \times (S^1 = \mathbb{R}/\mathbb{Z}) \rightarrow \mathbb{R}$, there is an associated time dependent Hamiltonian vector field X_t , $0 \leq t \leq 1$, defined by

$$(2.1) \quad \omega(X_t, \cdot) = -dH_t(\cdot).$$

The vector field X_t generates a path $\gamma : [0, 1] \rightarrow \text{Diff}(M)$, starting at id . Given such a path γ , its end point $\gamma(1)$ is called a Hamiltonian symplectomorphism. The space of Hamiltonian symplectomorphisms forms a group, denoted by $\text{Ham}(M, \omega)$. In particular the path γ above lies in $\text{Ham}(M, \omega)$. It is well-known that any smooth path γ in $\text{Ham}(M, \omega)$ with $\gamma(0) = \text{id}$ arises in this way (is generated by $H : M \times [0, 1] \rightarrow \mathbb{R}$ as above).

2.2. Floer chain complex. Classically, the generators of the Hamiltonian Floer chain complex associated to

$$H : M \times S^1 \rightarrow \mathbb{R},$$

are pairs $(o, \bar{o})$, with o a time 1 periodic orbit of the Hamiltonian flow generated by H , and $\bar{o}$ a homotopy class of a disk bounding the orbit. The function H determines

a Hamiltonian connection $\mathcal{A}_H$ on the bundle $M \times S^1 \rightarrow S^1$. The horizontal spaces for $\mathcal{A}_H$ are the $\tilde{\Omega}_H$ orthogonal spaces to the vertical tangent spaces of $M \times S^1$, where

$$\tilde{\Omega}_H = \omega - d(Hd\theta).$$

Notation 2.1. *If H generates p then we may also denote $\mathcal{A}_H$ by $\mathcal{A}_p$.*

The horizontal (a.k.a flat) sections for $\mathcal{A}_H$ correspond to periodic orbits of H , in the obvious way. The homotopy classes $\bar{o}$, induce homotopy classes of bounding disks in $M \times D^2$ of the corresponding flat sections. The connection $\mathcal{A}_H$ induces an obvious $\mathbb{R}$ -translation invariant connection on $M \times \mathbb{R} \times S^1$, trivial in the $\mathbb{R}$ direction, that we also call $\mathcal{A}_H$. For an $\mathbb{R}$ translation invariant family $\{j_\theta\}$ of almost complex structures on the vertical tangent bundle of $M \times \mathbb{R} \times S^1$, $\mathcal{A}_H$ induces an almost complex structure J_H on $M \times \mathbb{R} \times S^1$, by declaring $\mathcal{A}_H$ horizontal spaces to be J_H invariant, requiring that J_H coincides with $\{j_\theta\}$ on the vertical tangent bundle, and asking the projection to $(\mathbb{R} \times S^1, j_{st})$ be J_H holomorphic.

Assuming the pair $(H, \{j_\theta\})$ is regular, (which is a generic property in both variables in the so called semi-positive case), the differential in the classical Hamiltonian Floer chain complex is obtained via count (modulo natural action of $\mathbb{R}$) of J_H -holomorphic sections u of $M \times \mathbb{R} \times S^1 \rightarrow \mathbb{R} \times S^1$, whose projections to $M \times S^1$ are asymptotic in backward, forward time to generators $(o_-, \bar{o}_-)$, respectively $(o_+, \bar{o}_+)$ of $CF(H)$, such that $\bar{o}_- + [u] - \bar{o}_+ = 0 \in \pi_2(M \times S^2)$, with the obvious interpretation of this equation, and such that

$$CZ(o_+, \bar{o}_+) - CZ(o_-, \bar{o}_-) = 1.$$

We shall omit further details. The corresponding chain complex will be denoted by $CF(p, \{j_\theta\})$, if p is the path from id to $\phi \in \text{Ham}(M, \omega)$ generated by H or just $CF(p)$ if $\{j_\theta\}$ is implicit.

Grading does not play an important role in this note except in order to specify in which degree the fundamental class lies. We shall grade the Floer chain complex using Conley-Zehnder index so that for a C^2 small, Morse, time independent Hamiltonian $H : M \rightarrow \mathbb{R}$, the CZ index of a flat section of $\mathcal{A}_H$ corresponding to a local maximizer is $2n$.

3. PROOF OF THEOREM 1.2

Fix $b_0 \in B$. We consider the path Serre fibration $\mathcal{P}^B$ over B , with fiber $\mathcal{P}^B(b)$ over b the space of paths in B from b_0 to b . We shall always assume that our paths are constant near end points. While each fiber $\mathcal{P}^B(b)$ may not be contractible, its induced map into $\mathcal{P}(j(b))$: the space of all paths in $\text{Ham}(S^2, \omega)$ from $j(b_0)$ to $j(b)$, will be shown to be null-homotopic, as follows. Associated to each $p \in \mathcal{P}^B(b)$ we get a Hamiltonian fibration C_p over $\mathbb{CP}^1$, formed via the clutching construction using the loop $j_*p_0 \cdot j_*p^{-1}$, where $p_0 \in \mathcal{P}^B(b)$ is a certain reference path. The non degeneracy assumption is to construct a natural foliation of C_p by holomorphic sections. This foliation in turn determines a natural smooth trivialization of C_p . The construction of this foliation also uses an elementary application of Cerf theory, together with classical positivity of intersections ideas of Gromov-McDuff, particularly following [2]. Given this it is a matter of simple topology to deduce the main result.

Proposition 3.1. *The natural map*

$$i_* j_* : \mathcal{P}^B(b) \rightarrow \mathcal{P}(j(b))$$

is vanishing on all homotopy groups.

Proof. We may suppose that B is connected so that j maps into $\text{Ham}(S^2, 2k)$ for some $2k$. Fix $p_0 \in \mathcal{P}^B(b)$, so that the loop $f(p) = j_* p_0 \cdot j_* p^{-1}$ is contractible in $\mathcal{P}(j(b))$ for some and hence every p . Then for each $p \in \mathcal{P}^B(b)$ we get a Hamiltonian S^2 fibration over $S^1 \times \mathbb{R}$ that we call X_p by using $f(p)$ as a clutching map:

$$(3.1) \quad X_p = S^2 \times (-\infty, 0] \times S^1 \sqcup S^2 \times [0, \infty) \times S^1 / \sim,$$

for $\sim$ the equivalence relation under which

$$(x, (0, \theta)) \in S^2 \times \partial(-\infty, 0] \times S^1$$

is equivalent to

$$(f^{-1}(p)_\theta(x), (0, \theta)) \in S^2 \times \partial[0, \infty) \times S^1,$$

using the polar coordinates (r, θ) , $0 \leq \theta \leq 1$. Fix any path $p_{b_0} \in \mathcal{P}(j(b_0))$. This determines a translation invariant Hamiltonian connection over the ends $(-\infty, -1]$ and $[1, \infty)$ of X_p , which over $\{r\} \times S^1$, $r \leq -1$, $r \geq 1$ is the Hamiltonian connection with holonomy path generated by $H^{p_{b_0}}$ the generating function for p_{b_0} , and is trivial in the r direction. The paths p_0, p partially determine a connection $\mathcal{A}_{p_0, p}$ extending the above connection over all $\mathbb{R} \times S^1$, so that for $r \in [-1, 0]$ the flat sections of $\mathcal{A}_{p_0, p}$ over $\{r\} \times S^1$ correspond to fixed points of $j_* p_0(1+r)$, and for $r \in [0, 1]$ the flat sections of $\mathcal{A}_{p_0, p}$ over $\{r\} \times S^1$ correspond to fixed points of $j_* p(r-1)$. In the r direction $\mathcal{A}_{p_0, p}$ is specified arbitrarily.

By assumption for each p the pair $(X_p|_{\{r\} \times S^1}, \mathcal{A}_{p_0, p})$ is Floer non-degenerate for all r , meaning the holonomy map of the connection over S^1 has non-degenerate fixed points. Consequently we have $2k$ smooth sections $\{\sigma_i\}$ of X_p whose restrictions over $\{r\} \times S^1$ for each r are flat sections of $\mathcal{A}_{p_0, p}$, corresponding to the fixed points of the associated holonomy map. We shall call these *spectral sections*. The asymptotic constraint at $\pm\infty$ for each σ_i will be denoted by $o_{i, \pm}$. Let us label the generators of $CF(p_{b_0})$ by $\tilde{o}_{i, -} = (o_{i, -}, \tilde{o}_{i, -})$. Then we may further specify that the asymptotic constraints of each σ_i are $\tilde{o}_{i, \pm}$, i.e. taking homotopy classes of bonding disks into account, with $\tilde{o}_{i, +}$ also in $CF(p_{b_0})$, this makes sense because each loop $f(p)$ is contractible by assumptions, so that starting with a homotopy class of a bounding disk for $o_{i, -}$ in $M \times D^2$, and trivializing X_p relative to the ends, σ_i induces a homotopy class of a bounding disk for $o_{i, +}$ and it is immediate that resulting CZ index of $\tilde{o}_{i, +}$ coincides with the CZ index for $\tilde{o}_{i, -}$. The CZ index 2 generators will be denoted by $\{\tilde{o}_{j, \pm}\}$ (with $\{o_{j, \pm}\}$ a subset of $\{o_{i, \pm}\}$).

Let $\mathcal{F}(p)$ denote the space of Hamiltonian connections on X_p coinciding with $\mathcal{A}_{p_0, p}$ over all the slices $\{r\} \times S^1$ and for which all the spectral sections are flat. This is an affine (and non empty) space and hence contractible. A Hamiltonian connection gives rise to an almost complex structure as described in the preliminaries.

The total space of the family $\mathcal{F}(p)$ fibers over $\mathcal{P}^B(p)$ and this is clearly a Serre fibration. The fiber is contractible as previously observed so there are sections. Let $g : S^m \rightarrow \mathcal{P}^B(b)$ be a continuous map. Pullback $\{\mathcal{F}(p)\}$ by g and take any section $\{\mathcal{A}_s\}$, $s \in S^m$ and let $\{J_s\}$ denote the corresponding family of almost complex structures on $\{X_s\} = \{X_{g(s)}\}$. Cap off both ends of X_s via gluing, for a

sufficiently small gluing parameter (to be described shortly) with $(M \times \mathbb{C}, \mathcal{A}_{p_{b_0}}^{cap})$ for $\mathcal{A}_{p_{b_0}}^{cap}$ the Hamiltonian connection with holonomy path p_{b_0} over radius r circles in $\mathbb{C}$ for r sufficiently large, and trivial in the radial direction for r large. We call the resulting glued structure C_s , which is a Hamiltonian S^2 fibration over $\mathbb{CP}^1$, and the resulting Hamiltonian connection on C_s will be called $\mathcal{A}_s^{cap}$.

We may suppose that the path p_0 was chosen so that the induced Hamiltonian connection $\mathcal{A}_{p_{b_0}}$, is regular that is so that $CF(p_{b_0})$ is defined. We likewise choose $\tilde{\mathcal{A}}_{p_{b_0}}$ extending $\mathcal{A}_{p_{b_0}}$ in the natural sense, so that the associated moduli spaces

$$\mathcal{M}(\tilde{\mathcal{A}}_{p_{b_0}}, \tilde{o}_{j,\pm})$$

of $J(\tilde{\mathcal{A}}_{p_{b_0}})$ -holomorphic sections of $M \times \mathbb{C}$ asymptotic to $\tilde{o}_{j,\pm}$ are regular. As a remark we note that the Fredholm index will be 0 for elements of $\mathcal{M}(\tilde{\mathcal{A}}_{p_{b_0}}, \tilde{o}_{j,-})$ and Fredholm index 2 for $\mathcal{M}(\tilde{\mathcal{A}}_{p_{b_0}}, \tilde{o}_{j,+})$ as we must reverse the orientation on $\mathbb{C}$ for the $+$ end to do the gluing, which affects how the Fredholm index is determined by Atiyah-Singer index theorem via the Conley-Zehnder index. This is as expected of course since we must get 2 dimensional moduli space after gluing. We shall not elaborate as this is a rather elementary technical point and more of a remark to the reader.

Consequently we obtain a family $\{C_s, J_s^{cap}\}$ of Hamiltonian fibrations over $\mathbb{CP}^1$ each of which is Hamiltonian equivalent to the trivial bundle $S^2 \times \mathbb{CP}^1$, by the assumption that $i \circ j$ vanishes on π_1 .

Lemma 3.2. *If the gluing parameter was taken to be sufficiently small the moduli space $\overline{\mathcal{M}}(s, J_s^{cap})$ of J_s^{cap} stable holomorphic sections of C_s , in the class of a constant section $[const]$ contains no nodal curves. Where J_s^{cap} denotes the almost complex structure induced by $\mathcal{A}_s^{cap}$.*

Proof. Let

$$\sum_j c_j \tilde{o}_{j,-},$$

represent a fundamental class in $CF_2(p_{b_0})$. Then for some j' with $c_{j'} \neq 0$ there must be an element σ'_- of $\mathcal{M}(\mathcal{A}_{p_{b_0}}, \tilde{o}_{j',-})$ since the PSS map from quantum homology to Floer homology is obtained by count of elements of these moduli spaces and so must hit this fundamental chain. The chain

$$\sum_j c_j \tilde{o}_{j,+},$$

must also represent the fundamental class in $CF_2(p_{b_0})$. This fact is an extremely special case of Cerf-Floer continuation map theory, which in this case is without bifurcations, and follows by classical induced cobordisms of moduli spaces technique. If however the reader is curious about the general framework for Cerf-Floer theory with bifurcations, we refer to [4] or Oh [7] for a more recent treatment. So there likewise must be an element σ'_+ in the moduli space $\mathcal{M}(\mathcal{A}_{p_{b_0}}, \tilde{o}_{j',+})$.

Now, the spectral section $\sigma_{j'}$ is regular by automatic transversality, [6, Appendix C], consequently if the gluing parameter was small we may analytically glue σ'_- , σ'_+ and $\sigma_{j'}$ to get a smooth J_p^{cap} holomorphic $[const]$ class section σ^{cap} of C_p . Finally, if there are nodal curves in our moduli space $\overline{\mathcal{M}}(p, J_p^{cap})$ then taking the principal component of the corresponding nodal section, we would have a smooth

holomorphic section σ' in class $[const] + A$ for some spherical fiber class A with $c_1(A) < 0$. Consequently σ' would have negative intersection number with σ^{cap} , which is impossible by positivity of intersections as σ^{cap} is embedded, see [6, Section 2.6]. $\square$

Lemma 3.3. $\overline{\mathcal{M}}(s, J_s^{cap})$ is also regular, (in particular has the expected dimension).

Proof. By automatic transversality the top stratum is regular, as since all elements are in class $[const]$, the associated real linear CR operator is automatically surjective, see for example [6, Appendix C]. But we just showed that there are no nodal curves and hence no other strata. $\square$

By regularity and compactness the map

$$ev_s : \overline{\mathcal{M}}(s, J_s^{cap}) \rightarrow \pi_s^{-1}(z_0) \simeq S^2,$$

taking a section to its value over some $z_0 \in \mathbb{C}$ is a smooth degree 1 map. Together with positivity of intersections we get that the sections of $\overline{\mathcal{M}}(s, J_s^{cap})$ induce a canonical smooth foliation of C_s , with all leaves diffeomorphic to $\mathbb{CP}^1$, and as they are sections we get a canonical smooth trivialization

$$C_s \xrightarrow{tr_s} S^2 \times \mathbb{CP}^1,$$

by identifying all fibers with a fixed fiber over $0 \in \mathbb{CP}^1$ using the leaves of the foliation.

We get a bundle $P_g \rightarrow S^m$ with fiber over s : C_s and with structure group reducible to the group of Hamiltonian bundle automorphisms of $S^2 \times \mathbb{CP}^1$ fixing the fiber over 0, with 0 identified with the origin of $\mathbb{C}$ before gluing in the cap $S^2 \times \mathbb{C}$ at the $-\infty$ end of X_s . In other words this is the group:

$$(3.2) \quad \Omega_{id}^2 \text{Ham}(S^2, \omega),$$

with Ω^2 denoting the based spherical mapping space. We drop the subscript id from now on. We have an analogous group

$$\Omega^2 \text{Diff}(S^2)$$

consisting of smooth orientation preserving bundle automorphisms fixing the fiber over 0. By the argument above the bundle P_g is trivializable as a structure group $\Omega^2 \text{Diff}(S^2)$ bundle. The inclusion

$$\Omega^2 \text{Ham}(S^2, \omega) \rightarrow \Omega^2 \text{Diff}(S^2),$$

is a homotopy equivalence by the classical theorem of Smale, [8], for this particular version we can also use Moser argument. Consequently P_g is trivializable as a structure group $\Omega^2 \text{Ham}(S^2, \omega)$ bundle.

By construction P_g is equivalent to the pullback by the sequence

$$S^m \xrightarrow{f \circ j \circ g} \Omega_{j(b_0)} \text{Ham}^{ndg}(S^2) \xrightarrow{\tilde{i}} \Omega_{j(b_0)} \text{Ham}(S^2, \omega) \rightarrow \Omega_{id} \text{Ham}(S^2, \omega) \rightarrow \Omega^2 B\text{Ham}(S^2, \omega)$$

of the universal structure group $\Omega^2 \text{Ham}(S^2, \omega)$ bundle over $\Omega^2 B\text{Ham}(S^2, \omega)$. The last pair of maps are homotopy equivalences, so if P is trivial as a structure group $\Omega^2 \text{Ham}(S^2, \omega)$ bundle the map $\tilde{i} \circ (f \circ j \circ g)$ is null-homotopic. It immediately follows that $i_* j_*$ is vanishing on all homotopy groups. $\square$

4. ACKNOWLEDGEMENTS

The paper was completed at ICMAT Madrid, where the author is a research fellow, with funding from Severo Ochoa grant and Viktor Ginzburg laboratory. Michael Usher for reading of an early draft and pointing out a gap, to Dusa McDuff for interest, and much help in clarifying the paper. Viktor Ginzburg for interest and comments.

REFERENCES

- [1] M. AKVELD AND D. SALAMON, *Loops of Lagrangian submanifolds and pseudoholomorphic discs.*, Geom. Funct. Anal., 11 (2001), pp. 609–650.
- [2] M. GROMOV, *Pseudo holomorphic curves in symplectic manifolds.*, Invent. Math., 82 (1985), pp. 307–347.
- [3] S. HU AND F. LALONDE, *A relative Seidel morphism and the Albers map.*, Trans. Am. Math. Soc., 362 (2010), pp. 1135–1168.
- [4] Y. LEE, *Reidemeister torsion in Floer-Novikov theory and counting pseudo-holomorphic tori. I.*, J. Symplectic Geom., 3 (2005), pp. 221–311.
- [5] D. MCDUFF, *Configuration spaces of positive and negative particles.*, Topology, 14 (1975), pp. 91–107.
- [6] D. MCDUFF AND D. SALAMON, *J-holomorphic curves and symplectic topology*, no. 52 in American Math. Society Colloquium Publ., Amer. Math. Soc., 2004.
- [7] Y.-G. OH, *Floer mini-max theory, the Cerf diagram, and the spectral invariants.*, J. Korean Math. Soc., 46 (2009), pp. 363–447.
- [8] S. SMALE, *Diffeomorphisms of the 2-sphere.*, Proc. Am. Math. Soc., 10 (1959), pp. 621–626.

ICMAT, MADRID

E-mail address: yasha.savelyev@gmail.com